



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Foreign and insular statistical reports of countries and cities—Yearly and monthly.*

BAHAMAS—*Dunmore Town*.—Two weeks ended January 25, 1901. Estimated population, 1,472. No deaths and no contagious diseases.

*Governors Harbor*.—Two weeks ended February 2, 1901. Estimated population, 1,500. No deaths and no contagious diseases.

*Green Turtle Cay—Abaco*.—Two weeks ended January 31, 1901. Estimated population, 3,900. No deaths and no contagious diseases.

*Inagua*.—Four weeks ended January 28, 1901. Estimated population, 1,200. No deaths.

*Nassau*.—Two weeks ended February 6, 1901. Estimated population, 12,000. No deaths and no contagious diseases reported.

BRAZIL—*Rio de Janeiro*.—Two weeks ended December 15, 1900. Estimated population, 779,000. Total number of deaths, 563, including diphtheria, 2; enteric fever, 2; beriberi, 3; smallpox, 27; plague, 7, and 114 from tuberculosis.

CANADA—*Manitoba—Winnipeg*.—Estimated population, 25,642. Total number of deaths not reported. Two deaths from diphtheria, 1 death from enteric fever, 1 from scarlet fever, and 2 deaths from tuberculosis reported.

QUEBEC—*Sherbrooke*.—Month of January, 1901. Estimated population, 11,100. Total number of deaths, 24, including diphtheria, 1; enteric fever, 1, and 10 from scarlet fever.

CHILE—*Antofagasta*.—Month of December, 1900. Estimated population, 18,883. Total number of deaths, 62, including enteric fever, 1; whooping cough 9, and 5 from phthisis pulmonalis.

FRANCE—*Rouen*.—Month of December, 1900. Estimated population, 112,657. Total number of deaths, 203, including enteric fever 6, and 33 from tuberculosis.

GERMANY—*Glauchau*.—Month of November, 1900. Estimated population, 28,405. Total number of deaths 26, including 4 from tuberculosis.

*Hanover*.—Month of November, 1900. Estimated population, 235,572. Total number of deaths, 264, including diphtheria, 7; enteric fever, 2, and 28 from phthisis pulmonalis.

*Magdeburg*.—Month of October, 1900. Estimated population, 229,944. Total number of deaths, 345, including diphtheria, 5; enteric fever, 1; measles, 8; whooping cough, 3, and 30 from phthisis pulmonalis.

*Weimar*.—Month of December, 1900. Estimated population, 29,633. Total number of deaths, 43. No contagious diseases reported.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended January 19, 1901, correspond to an annual rate of 19.1 a thousand of the aggregate population, which is estimated at 11,788,099. The highest rate was recorded in Preston, viz, 24.7, and the lowest in Croydon, viz, 13.9.

*London*.—One thousand six hundred and sixty-six deaths were registered during the week, including measles, 36; scarlet fever, 9;

diphtheria, .32; whooping cough, 38; enteric fever, 13, and diarrhea and dysentery, 15. The deaths from all causes correspond to an annual rate of 18.7 a thousand. In Greater London 2,234 deaths were registered, corresponding to an annual rate of 17.2 a thousand of the population. In the "outer ring" the deaths included 16 from diphtheria, 10 from measles, and 12 from whooping cough.

*Ireland.*—The average annual death rate represented by the deaths registered during the week ended January 19, 1901, in the 22 principal town districts of Ireland was 26.0 a thousand of the population, which is estimated at 1,058,722. The lowest rate was recorded in Armagh, viz, 7.1, and the highest in Carrickfergus, viz, 40.9 a thousand. In Dublin and suburbs 209 deaths were registered, including enteric fever, 3, and 12 from whooping cough.

*Scotland.*—The deaths registered in 8 principal towns during the week ended January 19, 1901, correspond to an annual rate of 24.0 a thousand of the population, which is estimated at 1,630,155. The lowest mortality was recorded in Leith, viz, 15.0, and the highest in Aberdeen, viz, 29.3 a thousand. The aggregate number of deaths registered from all causes was 752, including diphtheria, 6; measles, 23; scarlet fever, 2; smallpox, 3, and 61 from whooping cough.

*JAMAICA—Port Antonio.*—Two weeks ended January 19, 1901. Population not reported. Number of deaths not reported. The health of the port is good.

*JAVA—Batavia.*—Three weeks ended December 29, 1900. Estimated population, 150,000. Number of deaths not reported. No contagious or epidemic diseases.

*MOROCCO—Tangier.*—Five weeks ended January 19, 1901. Estimated population, 40,000. No deaths and no contagious diseases reported.

*ST. HELENA.*—Two weeks ended December 29, 1900. Estimated population, 1,470. Total number of deaths, 3. No contagious diseases.

*SPAIN—Cadiz.*—Month of December, 1900. Estimated population, 67,987. Total number of deaths, 200, including diphtheria 1, and 2 from enteric fever.

*Malaga.*—Two weeks ended December 31, 1900. Estimated population, 130,000. Total number of deaths, 198, including enteric fever 1, and 1 from leprosy.

*Valencia.*—Two weeks ended January 13, 1901. Estimated population, 203,985. Total number of deaths, 203, including 1 from enteric fever.